The Dirac Equation Derived using 4-Vectors

It can be shown that spin comes from the Poincaré Symmetry of SR, not from a QM axiom. Writing a 4-SpinMomentum then leads naturally to the Dirac Equation.

First, we will do a little pure mathematics: Let { $(a^2 - b^2) = (a+b)(a-b) = c^2$ }

We can multiply by an arbitrary factor (xy): $(a^2 - b^2)(xy) = (a+b)(a-b)(xy) = c^2(xy)$

If we impose the following extra constraint: (a+b)x = (cy) (a-b)y = (cx)

Then the separated equations are still true when multiplied together: $(a+b)x * (a-b)y = (cy) * (cx) \rightarrow (a+b)(a-b)(xy) = c^{2}(xy)$

Now add and subtract the separated equations: (a+b)x + (a-b)y = (cy) + (cx)(a+b)x - (a-b)y = (cy) - (cx)

Gather terms in $\{a,b,c\}$: a(x+y) + b(x-y) = c(x+y)a(x-y) + b(x+y) = -c(x-y)

Let X=(x+y) and Y=(y-x)=-(x-y), just a change in variable names aX - bY = cX-aY + bX = cY

Rearrange: aX - bY = cXbX - aY = cY

Putting into matrix form:

 $\begin{array}{l} \begin{bmatrix} a - b \end{bmatrix} & \begin{bmatrix} X \\ b - a \end{bmatrix} & \begin{bmatrix} Y \end{bmatrix} & = \begin{bmatrix} c & 0 \end{bmatrix} & \begin{bmatrix} X \\ 0 & c \end{bmatrix} & \begin{bmatrix} Y \end{bmatrix} \\ or \\ Putting into suggestive matrix form... \\ \begin{pmatrix} \begin{bmatrix} 1 & 0 \\ 0 - 1 \end{bmatrix} & a + \begin{bmatrix} 0 - 1 \\ 1 & 0 \end{bmatrix} & \begin{bmatrix} X \\ 0 \end{bmatrix} = c \begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} X \\ 0 & 1 \end{bmatrix} \begin{bmatrix} Y \end{bmatrix}$

And again, to confirm that this matches the original equation:

aX - bY = cX bX - aY = cYor (a-c)X = (b)Y(b)X = (a+c)Y Multiply the terms: (a-c)X(a+c)Y = (b)X(b)Y(a-c)(a+c)XY = (b)(b)XY

 $(a^2 - c^2)XY = (b^2)XY$ $(a^2 - c^2) = b^2$ $(a^2 - b^2) = c^2$

So, mathematically:

 $(a^2 - b^2) = c^2$ is the defining equation, which holds equivalently for:

 $\{ (a^2 - b^2)(xy) = (c^2)(xy) \}$ or $(\begin{bmatrix} 1 & 0 \end{bmatrix} \\ (\begin{bmatrix} 0 & -1 \end{bmatrix} \\ a + \begin{bmatrix} 0 & -1 \end{bmatrix} \\ \begin{bmatrix} 1 & 0 \end{bmatrix} \\ b \end{bmatrix} [X] = c \begin{bmatrix} 1 & 0 \end{bmatrix} \\ \begin{bmatrix} 0 & 1 \end{bmatrix} \\ [Y]$

It is only the nature of $\{x,y\}$ and $\{X,Y\}$ that are different.

<u>Pauli Matrices</u> are a set of dimensionless 2x2 matrices that may be used in totally classical contexts. In this regard, they are not a quantum postulate, but one of Poincaré Invariance.

$$\sigma^{t} = \sigma^{0} = \mathbf{I}_{2} = \begin{bmatrix} 1 & 0 \end{bmatrix} \quad \sigma^{x} = \sigma^{1} = \begin{bmatrix} 0 & 1 \end{bmatrix} \quad \sigma^{y} = \sigma^{2} = \begin{bmatrix} 0 & -i \end{bmatrix} \quad \sigma^{z} = \sigma^{3} = \begin{bmatrix} 1 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 1 \end{bmatrix} \quad \begin{bmatrix} 1 & 0 \end{bmatrix} \quad \begin{bmatrix} 1 & 0 \end{bmatrix} \quad \begin{bmatrix} 0 & -1 \end{bmatrix}$$

The Pauli Spin Matrices can be written in Tensor notation, with each component itself a dimensionless 2x2 matrix $\Sigma = \Sigma^{\mu}{}_{v} = \text{Diag}[\sigma^{0}, \sigma^{1}, \sigma^{2}, \sigma^{3}] = \text{Diag}[\sigma^{0}, \sigma] \rightarrow \text{Diag}[\sigma^{i}, \sigma^{x}, \sigma^{y}, \sigma^{z}]$

In <u>classical mechanics</u>, Pauli matrices are useful in the context of the Cayley-Klein parameters. The 3D matrix *P* corresponding to the 3-position **x** of a point in space is defined in terms of the above Pauli vector matrix, $P = \mathbf{x} \cdot \mathbf{\sigma} = \mathbf{x} \cdot \mathbf{\sigma}^{x} + \mathbf{y} \cdot \mathbf{\sigma}^{y} + \mathbf{z} \cdot \mathbf{\sigma}^{z}$

The 4D matrix *P* corresponding to the 4-Position X of a point in spacetime is defined in terms of the above Pauli vector matrix, $P = X \cdot \Sigma = \operatorname{ct} \sigma^{t} + x \sigma^{x} + y \sigma^{y} + z \sigma^{z}$

Suppose now that the vector $\mathbf{x} = (x_1, x_2, x_3)$ is rotated around an axis with unit vector $\mathbf{n} = (n_1, n_2, n_3)$ through an angle θ .

The transformation matrix $\mathbf{R}(\theta)$ for rotations about an axis through an angle θ may be written in terms of Pauli matrices and the unit matrix. These can be gathered together in tensor notation with the 2D unit matrix = $\mathbf{I}_2 = \sigma^0$.

 $\boldsymbol{R}_{x}(\theta) = e^{-i\theta x/2} = \sigma^{0} \cos(\theta/2) - i \sigma^{x} \sin(\theta/2) = \begin{bmatrix} \cos(\theta/2) & -i \sin(\theta/2) \end{bmatrix} \\ \begin{bmatrix} -i \sin(\theta/2) & \cos(\theta/2) \end{bmatrix}$

 $\boldsymbol{R}_{y}(\theta) = e^{-i\theta y/2} = \sigma^{0} \cos(\theta/2) - i \sigma^{y} \sin(\theta/2) = \begin{bmatrix} \cos(\theta/2) & -\sin(\theta/2) \end{bmatrix} \\ \begin{bmatrix} \sin(\theta/2) & \cos(\theta/2) \end{bmatrix}$

 $\boldsymbol{R}_{z}(\theta) = e^{-i\theta z/2} = \sigma^{0} \cos(\theta/2) - i \sigma^{z} \sin(\theta/2) = \begin{bmatrix} \cos(\theta/2) - i \sin(\theta/2) & 0 \\ 0 & \cos(\theta/2) + i \sin(\theta/2) \end{bmatrix} = e^{-i(\theta/2)} 0 = e^{-i(\theta/2)} e^{-i(\theta/$

For an angle θ about an arbitrary axis **n**

$$\boldsymbol{R}_{\boldsymbol{n}}(\theta) = e^{-i\theta n/2} = \sigma^{0} \cos(\theta/2) - i (\mathbf{n} \cdot \boldsymbol{\sigma}) \sin(\theta/2) = \begin{bmatrix} \cos(\theta/2) - i \sin(\theta/2)n_{3} & -\sin(\theta/2)(n_{2}+in_{1}) \end{bmatrix} \\ \begin{bmatrix} \sin(\theta/2)(n_{2}-in_{1}) & \cos(\theta/2)+i\sin(\theta/2)n_{3} \end{bmatrix}$$

All of the relativistic wave equations can be derived from a common source: The relativistic mass-energy relation, including spin, in an Electromagnetic (EM) field. Note that this formalism fits well with the Stern-Gerlach experiment.

4-Momentum P = (E/c,p)

4-Momentum including Spin $\mathbf{P}_{\mathbf{s}} = \boldsymbol{\Sigma} \cdot \mathbf{P} = \Sigma^{\mu} {}_{\mathbf{v}} \mathbf{P}^{\nu} = \eta_{\alpha\beta} \Sigma^{\mu\alpha} \mathbf{P}^{\beta} = \mathbf{P} \mathbf{s}^{\mu}$ $\Sigma^{\mu} {}_{\mathbf{v}}$ is the 4D (1,1)-Pauli Spin-Matrix Tensor = Diag[$\sigma^{0}, \boldsymbol{\sigma}$] $\Sigma^{\mu\nu}$ is the 4D (2,0)-Pauli Spin-Matrix Tensor = Diag[$\sigma^{0}, \boldsymbol{\sigma}$]

 $\mathbf{P}_{\mathbf{s}} = \text{Diag}[\sigma^{0}, -\boldsymbol{\sigma}] \cdot \mathbf{P} = \text{Diag}[\sigma^{0}, -\boldsymbol{\sigma}] \cdot (E/c, \mathbf{p}) = (\sigma^{0}E/c, \boldsymbol{\sigma} \cdot \mathbf{p})$

 $\mathbf{P}_{\mathbf{S}} = (\mathbf{p}_{\mathbf{S}}^{0}, \mathbf{p}_{\mathbf{S}}) = (\sigma^{0} \mathbf{E} / \mathbf{c}, \boldsymbol{\sigma} \cdot \mathbf{p})$

with σ^0 as an identity matrix I of appropriate spin dimension and σ is the Pauli Spin Matrix Vector

4-Momentum inc. Spin in External Field $P_T = (H/c,p_T) = (E_T/c,p_T)$ with: $H = E_T =$ Hamiltonian = Total Energy Of System p_T = Total 3-momentum Of System

4-TotalMomentum $\mathbf{P}_{T} = \mathbf{P} + q\mathbf{A}$ 4-Momentum $\mathbf{P} = \mathbf{P}_{T} - q\mathbf{A}$ 4-MomentumIncSpin $\mathbf{P}_{s} = (\mathbf{p}_{s}^{0}, \mathbf{p}_{s}) = (\sigma^{0}\mathbf{E}/c, \boldsymbol{\sigma} \cdot \mathbf{p}) = (\sigma^{0}(\mathbf{E}_{T}/c - q\boldsymbol{\phi}/c), \boldsymbol{\sigma} \cdot (\mathbf{p}_{T} - q\mathbf{a}))$: Note each component is a 2x2 matrix $\mathbf{P}_{s} \cdot \mathbf{P}_{s} = (\mathbf{p}_{s}^{0})^{2} - (\mathbf{p}_{s})^{2} = [\sigma^{0}(\mathbf{E}/c)]^{2} - [\boldsymbol{\sigma} \cdot (\mathbf{p}_{T} - q\boldsymbol{\phi}/c)]^{2} - [\boldsymbol{\sigma} \cdot (\mathbf{p}_{T} - q\mathbf{a})]^{2} = (\mathbf{m}_{o}c)^{2} = (\mathbf{E}_{o}/c)^{2}$

The 4-TotalMomentum (inc. External Field Minimal-Coupling and Spin) $\mathbf{P}_{\mathbf{s}} = \boldsymbol{\Sigma} \cdot \mathbf{P} = \boldsymbol{\Sigma} \cdot (\mathbf{P}_{\mathbf{T}} - q\mathbf{A}) = [\sigma^{0}(\mathbf{E}_{\mathbf{T}}/\mathbf{c} - q\phi/\mathbf{c}), \boldsymbol{\sigma} \cdot (\mathbf{p}_{\mathbf{T}} - q\mathbf{a})]$ with $\boldsymbol{\Sigma} = \Sigma^{\mu\nu}$ as the Pauli Spin Matrices, and taking the Einstein summation gives the σ^{0} and $\boldsymbol{\sigma}$

$$\begin{split} \mathbf{P}_{\mathbf{s}} \cdot \mathbf{P}_{\mathbf{s}} &= (\boldsymbol{\Sigma} \cdot \mathbf{P})^2 = [\boldsymbol{\Sigma} \cdot (\mathbf{P}_{\mathbf{T}} - q\mathbf{A})]^2 = [\sigma^0 (\mathbf{E}_{\mathrm{T}}/\mathbf{c} - q\phi/\mathbf{c})]^2 - [\boldsymbol{\sigma} \cdot (\mathbf{p}_{\mathrm{T}} - q\mathbf{a})]^2 = (\mathbf{m}_{\mathrm{o}}\mathbf{c})^2 \\ &(\boldsymbol{\Sigma} \cdot \mathbf{P})^2 = (\mathbf{m}_{\mathrm{o}}\mathbf{c})^2 \\ &(\boldsymbol{\Sigma} \cdot \partial)^2 = -(\mathbf{m}_{\mathrm{o}}/\hbar)^2 \\ &(\boldsymbol{\Sigma} \cdot \partial)^2 + (\mathbf{m}_{\mathrm{o}}\mathbf{c}/\hbar)^2 = 0 \\ &(\boldsymbol{\Sigma} \cdot (\mathbf{D} - (i/h)q\mathbf{A}))^2 + (\mathbf{m}_{\mathrm{o}}\mathbf{c}/\hbar)^2 = 0 \end{split}$$

Now, to prove that this "Relativistic Pauli" Energy-Momentum equation can give the Dirac equation $\begin{aligned} \mathbf{Ps} \cdot \mathbf{Ps} &= [\sigma^0(\mathbf{E}_T/\mathbf{c} - \mathbf{q}\phi/\mathbf{c})]^2 - [\boldsymbol{\sigma} \cdot (\mathbf{p_T} - \mathbf{qa})]^2 = (\mathbf{p_s}^0)^2 - (\mathbf{p_s})^2 = (\mathbf{m_oc})^2 = (\mathbf{E_o/c})^2 \\ \mathbf{Ps} \cdot \mathbf{Ps} &= I(\mathbf{E}_T/\mathbf{c} - \mathbf{q}\phi/\mathbf{c})]^2 - [\boldsymbol{\sigma} \cdot (\mathbf{p_T} - \mathbf{qa})]^2 = (\mathbf{p_s}^0)^2 - (\mathbf{p_s})^2 = (\mathbf{m_oc})^2 = (\mathbf{E_o/c})^2 \\ \mathbf{Ps} \cdot \mathbf{Ps} &= (\mathbf{p_s}^0)^2 - (\mathbf{p_s})^2 = (\mathbf{p_s}^0 + \mathbf{p_s}) (\mathbf{p_s}^0 - \mathbf{p_s}) = (\mathbf{m_oc})^2 \\ \{(\mathbf{p_s}^0)^2 - (\mathbf{p_s})^2\}(\mathbf{xy}) = (\mathbf{m_oc})^2(\mathbf{xy})
\end{aligned}$

From our math proof above, this is equivalent to:

$$\begin{array}{cccc} ([1 & 0] & & & [0 - 1] & \\ ([0 - 1] & & p_s^0 + & & [1 & 0] & & p_s \end{array} \begin{array}{c})[X] & = (m_o c) I_2 & & [X] \\)[Y] & & & [Y] \end{array}$$

or

$$\begin{array}{cccc} ([1 & 0] & & & & [0 & -1] & \\ ([0 & -1] & & \sigma^0 p^0 + & & & [1 & 0] & & \sigma \cdot p & &) [X] & = (m_o c) I_2 & & & [X] \\ \end{array}$$

Putting into highly suggestive matrix form...

$$\begin{array}{cccc} ([\sigma^0 & 0] & & \\ ([0 & -\sigma^0] & & p^0 + & \begin{matrix} [0 & -\sigma] & \\ & & & \begin{matrix} [\sigma & 0] \end{matrix} & \cdot \textbf{p} \end{matrix} \begin{array}{c})[X] & = (m_o c) I_2 & & \begin{matrix} [X] \\ & & & \end{matrix} \\ [Y] \end{array}$$

 $\label{eq:prod} \begin{array}{ll} \mbox{let Spinor } \Psi = & \begin{array}{c} [X] \\ [Y] \end{array} \quad \mbox{and note that } \sigma^0 = I_2 \end{array}$

this is equivalent to Dirac Gamma Matrices (in Dirac Basis)...

$$\begin{array}{cccc} ([I_2 & 0] & & & & [0 \ \textbf{-\sigma}] \\ ([0 \ \textbf{-I}_2] & p^0 + & & [\textbf{\sigma} & 0] & \textbf{\cdot p} \end{array} \begin{array}{c}) & \Psi = (m_o c) I_2 \Psi \\ \end{array}$$

$$\begin{split} (\gamma^0 p^0 \boldsymbol{\cdot} \boldsymbol{\gamma} \boldsymbol{\cdot} \boldsymbol{p}) \Psi &= (m_o c) I \Psi \\ (\boldsymbol{\Gamma} \boldsymbol{\cdot} \boldsymbol{P}) \Psi &= (m_o c) I \Psi \\ (\boldsymbol{\Gamma} \boldsymbol{\cdot} \boldsymbol{P}) &= (m_o c) \\ (\Gamma^\mu P_\mu) \Psi &= (m_o c) \Psi \\ i\hbar (\Gamma^\mu \partial_\mu) \Psi &= (m_o c) \Psi \\ \end{split}$$
The Dirac Relativistic Quantum Equation for spin 1/2 particles

~

To recap, the SR 4-Vectors have some very simple, fundamental, and Invariant Lorentz Scalar relations between one another:			
4-Position	$\mathbf{R} = \mathbf{R}^{\mu} = (\text{ct}, \mathbf{r}) \in \langle \text{Event} \rangle \in \langle \text{Time} \cdot \text{Space} \rangle$	\rightarrow (E _o /c ²) -	$\mathbf{P} = (\mathbf{E/c,p})$
4-Velocity	$\mathbf{U} = \mathbf{U}^{\mu} = \gamma(\mathbf{c}, \mathbf{u}) = (\mathbf{U} \cdot \partial) \mathbf{R} = (\mathbf{d}/\mathbf{d}\tau) \mathbf{R}$	/	
4-Momentum	$\mathbf{P} = P^{\mu} = (E/c, \mathbf{p}) = (m_o)\mathbf{U}$	$\mathbf{U} = \gamma(\mathbf{c}, \mathbf{u})$	$(\hbar)\uparrow\downarrow(1/\hbar)$
4-WaveVector	$\mathbf{K} = \mathbf{K}^{\mu} = (\boldsymbol{\omega}/\mathbf{c}, \mathbf{k}) = (1/\hbar)\mathbf{P}$	\	
4-Gradient	$\partial = \partial^{\mu} = (\partial_t / c, -\nabla) = (-i)\mathbf{K}$	$\rightarrow (\omega_{o}/c^{2})$ w	→ K= (<u>ω/c,k</u>)

.

Analysis of Dirac's Constant ($\hbar = h/2\pi$): Planck's Constant ($h = 2\pi\hbar$), and ($\underline{E_0}/\omega_0$) in the context of SRQM:

It is an empirical (observational) fact that the Lorentz Scalar Invariant $(E_0/\omega_0) = (\gamma E_0/\gamma \omega_0) = (E/\omega) => (\hbar)$ [J·s]/{rad} for all known experimental measurements. The SR 4D-Tensor rules show that one doesn't need a quantum axiom for this. (\hbar) is actually an empirically-measurable quantity, just like (c), (e), (G), (k_B), (μ_0), (ϵ_0) or the other fundamental constants, which are also 4D Lorentz Scalar Invariants. (\hbar) can be measured classically {without need of quantum axioms} from the photoelectric effect, from the inverse photoelectric effect, from LED's (injection electroluminescence), from Atomic Line Spectra (Bohr Atom), from the Duane-Hunt Law in Bremsstrahlung, from Electron Diffraction in crystals, from the Watt/Kibble-Balance, from the Sagnac Effect, from Incandescent Blackbody Intensity-Temperature Relations, from Compton Scattering, from the Stern-Gerlach experiment, etc.

For physics simulations of measurements of Dirac's : Planck's Constant (h : h), see: http://scirealm.org/Physics-PlanckConstantViaAtomicLineSpectra.html http://scirealm.org/Physics-PlanckConstantViaComptonScattering.html http://scirealm.org/Physics-PlanckConstantViaElectronDiffraction.html http://scirealm.org/Physics-PlanckConstantViaGravityInducedInterferometry.html http://scirealm.org/Physics-PlanckConstantViaIncandescence.html http://scirealm.org/Physics-PlanckConstantViaLEDs.html http://scirealm.org/Physics-PlanckConstantViaSagnacEffect.html http://scirealm.org/Physics-PlanckConstantViaSagnacEffect.html http://scirealm.org/Physics-PlanckConstantViaSternGerlach.html

```
(transitions of electron energy-levels)
(photon:electron collisions)
(electron:atomic crystal scattering)
(gravity interaction)
(temperature interaction)
(electron:photon interaction)
(rotation interaction)
(magnetic-field interaction)
```

 $\{E_o / \omega_o = E / \omega = \hbar\}$ implies the following SR (wave:particle-like $\int \cdot \cdot$) relation:

 $\mathbf{P} = \hbar \mathbf{K} = (\mathbf{E/c, p}) = \hbar(\omega/c, \mathbf{k}) = (\hbar \omega/c, \hbar \mathbf{k})$

The temporal part $\{E = \hbar\omega\}$ gives Einstein's photoelectric quantum "postulate". Again, emphasis: <u>derived</u> from SR. The spatial part $\{p = \hbar k\}$ gives de Broglie's matter-wave quantum "postulate". Again, emphasis: <u>derived</u> from SR.

$\partial = -i\mathbf{K} = (\partial_t/\mathbf{c}, -\nabla) = -i(\omega/\mathbf{c}, \mathbf{k})$

The temporal part $\{\partial_t = -i\omega\}$ or $\{\omega = i\partial_t\}$ gives temporal: frequency complex planewave change: operator The spatial part $\{\nabla = i\mathbf{k}\}$ or $\{\mathbf{k} = -i\nabla\}$ gives spatial: wavenumber complex planewave change: operator

The standard Schrödinger QM Relations derived from SR:

Again, we examine these SR 4-Vector relations derived above... and by simply combining them...

 $\mathbf{P} = \hbar \mathbf{K}$ (a relation which is entirely empirical, based on just SR arguments, shown above)

 $\mathbf{K} = i\partial$ (which is a relation for complex plane-waves, used in classical EM and elsewhere)

 $\mathbf{P} = i\hbar\partial = (\mathbf{E/c,p}) = i\hbar(\partial_t/c, -\nabla)$

The temporal part { $E = i\hbar\partial_t = i\hbar\partial/\partial t$ } gives unitary QM time evolution operator. Again, emphasis: <u>derived</u> from SR. The spatial part { $p = -i\hbar V$ } gives the QM momentum operator. Again, emphasis: <u>derived</u> from SR.

 $\begin{array}{l} \mbox{Relativistic version (Relativistic Pauli Equation)} \\ \mbox{Ps} = [\underline{\sigma}^0 (E_T/c - q\phi/c)]^2 - [\underline{\sigma} \cdot (p_T - qa)]^2 = (m_o c)^2 = (E_o/c)^2 \\ [\underline{\sigma}^0 (E_T/c - q\phi/c)]^2 - [\underline{\sigma} \cdot (p_T - qa)]^2 = (m_o c)^2 \\ [\underline{I}(E_T/c - q\phi/c)]^2 - [\underline{\sigma} \cdot (p_T - qa)]^2 - i[\underline{\sigma} \cdot ((p_T \ x - qa) + (-qa \ x \ p_T))] = (m_o c)^2 \\ [\underline{I}(E_T/c - q\phi/c)]^2 - [\underline{I} \cdot (p_T - qa)]^2 - i[\underline{\sigma} \cdot ihq((-\nabla_T \ x - a) + (-a \ x \ -\nabla_T))] = (m_o c)^2 \\ [\underline{I}(E_T/c - q\phi/c)]^2 - [\underline{I} \cdot (p_T - qa)]^2 - i[\underline{\sigma} \cdot ihq((-\nabla_T \ x - a) + (-a \ x \ -\nabla_T))] = (m_o c)^2 \\ [\underline{I}(E_T/c - q\phi/c)]^2 - [\underline{I} \cdot (p_T - qa)]^2 - i[\underline{\sigma} \cdot ihq(B)] = (m_o c)^2 \\ [\underline{I}(E_T/c - q\phi/c)]^2 - [\underline{I} \cdot (p_T - qa)]^2 + hq[\underline{\sigma} \cdot B] = (m_o c)^2 \\ [\underline{I}(E_T/c - q\phi/c)]^2 = [\underline{I} \cdot (p_T - qa)]^2 - hq[\underline{\sigma} \cdot B] = (m_o c)^2 \end{array}$

Non-relativistic version (Standard Pauli Equation) $[\underline{I}(E_T/c - q\phi/c)]^2 - [\underline{\sigma} \cdot (p_T - qa)]^2 = (m_oc)^2$ $[\underline{I}(E_T/c - q\phi/c)]^2 = [\underline{\sigma} \cdot (p_T - qa)]^2 + (m_oc)^2$ $[\underline{I}(E_T/c - q\phi/c)] = (\pm) \text{ Sqrt } [[\underline{\sigma} \cdot (p_T - qa)]^2 + (m_oc)^2]$ $[\underline{I}(E_T/c - q\phi/c)] = (\pm) \text{ Sqrt } [[\underline{\sigma} \cdot (p_T - qa)]^2 + (m_oc)^2]$ $[\underline{I}(E_T/c - q\phi/c)] \sim (\pm) [[\underline{\sigma} \cdot (p_T - qa)]^2 - (m_oc)]$ $[\underline{I}(E_T/c - q\phi/c)] \sim (\pm) [([(p_T - qa)]^2 - hq[\underline{\sigma} \cdot B])/(2m_oc) + (m_oc)]$ $[\underline{I}(E_T - q\phi)] \sim (\pm) [([(p_T - qa)]^2 - hq[\underline{\sigma} \cdot B])/(2m_o) + (m_oc^2)]$ $E_T \sim q\phi (\pm) [([(p_T - qa)]^2 - hq[\underline{\sigma} \cdot B])/(2m_o) + (m_oc^2)]$ where the - (hq[\underline{\sigma} \cdot B])/(2m_o) is the Stern-Gerlach term